



However, if renal failure becomes established, then renal replacement therapy (RRT) may be needed to maintain homoeostasis. While there are no clear guidelines with respect to the ideal mode or timing of RRT, we will discuss pros and cons of the various bedside options” Flood and Nichol (2018).

Abstract:

Acute renal failure is commonly encountered in the intensive care unit. It is associated with considerable morbidity and mortality. There are many possible aetiologies in the critically ill, including nephrotoxic agents, hypovolaemia and sepsis. While many classification systems for acute renal failure exist, the RIFLE (Risk, Injury, Failure, Loss, End-stage) criteria and the Acute Kidney Injury Network (AKIN) criteria are the most commonly utilized.

ReTweet if useful... Criteria to determine acute kidney injury in the critically ill patient
<https://ctt.ec/e9OaD+> @ivteam #ivteam

Click To Tweet

Many supportive therapies are employed to minimize the degree of renal injury once recognized, such as fluid resuscitation and maintenance of an adequate mean arterial pressure (with the use of inotropes in persistent hypotension despite fluid and treatment of the underlying aetiology). However, if renal failure becomes established, then renal

replacement therapy (RRT) may be needed to maintain homoeostasis. While there are no clear guidelines with respect to the ideal mode or timing of RRT, we will discuss pros and cons of the various bedside options.

Reference:

Flood, L. and Nichol, A. (2018) Acute kidney injury and the critically ill. *Anaesthesia and Intensive Care Medicine*. February 7th. .

DOI: <https://doi.org/10.1016/j.mpaic.2017.12.006>

Thank you to our partners for supporting IVTEAM

