



Our trial demonstrates that critically ill neurologic patients who require a central venous catheter have significantly lower odds of ultrasound-diagnosed CRLVT with placement of a CICVC as compared to a PICC” Fletcher et al (2016).

Abstract:

BACKGROUND: Observational studies suggest peripherally inserted central venous catheters (PICCs) are associated with a high risk of catheter-related large vein thrombosis (CRLVT) in critically ill neurologic patients. We evaluated the difference in thrombosis risk between PICCs and centrally inserted central venous catheters (CICVCs).

ReTweet if useful... PICC and CICVC associated upper extremity deep vein thrombosis
<http://ctt.ec/66yvf+> @ivteam #ivteam

Click To Tweet

METHODS: We conducted a pragmatic, randomized controlled trial of critically ill adult neurologic patients admitted to neurological and trauma critical care units at two level I trauma centers. Patients were randomized to receive either a PICC or CICVC and undergo active surveillance for CRLVT or death within 15 days of catheter placement.

RESULTS: In total, 39 subjects received a PICC and 41 received a CICVC between February 2012 and July 2015. The trial was stopped after enrollment of 80 subjects due to feasibility

affected by slow enrollment and funding. In the primary intention-to-treat analysis, 17 (43.6 %) subjects that received a PICC compared to 9 (22.0 %) that received a CICVC experienced the composite of CRLVT or death, with a risk difference of 21.6 % (95 % CI 1.57-41.71 %). Adjusted common odds ratio of CRLVT/death was significantly higher among subjects randomized to receive a PICC (adjusted OR 3.08; 95 % CI 1.1-8.65). The higher adjusted odds ratio was driven by risk of CRLVT, which was higher in those randomized to PICC compared to CICVC (adjusted OR 4.66; 95 % CI 1.3-16.76) due to increased large vein thrombosis without a reduction in proximal deep venous thrombosis.

CONCLUSIONS: Our trial demonstrates that critically ill neurologic patients who require a central venous catheter have significantly lower odds of ultrasound-diagnosed CRLVT with placement of a CICVC as compared to a PICC.

Reference:

Fletcher, J.J., Wilson, T.J., Rajajee, V., Stetler, W.R. Jr., Jacobs, T.L., Sheehan, K.M. and Brown, D.L. (2016) A Randomized Trial of Central Venous Catheter Type and Thrombosis in Critically Ill Neurologic Patients. *Neurocritical Care*. February 2nd. .

Thank you to our partners for supporting IVTEAM

