

ChloroPrep 1.5ml applicator

2% Chlorhexidine Gluconate w/v and 70% Isopropyl Alcohol v/v

Licensed, single use, sterile, ANTT™, latex free, skin antiseptics system

Pinch

Holding the barrel of the applicator, pinch the levers together, being careful not to touch the sponge.

Pinch the lever to release the solution. You will hear a 'pop' as the ampoule breaks.



Apply

Gently press the applicator against the skin until the solution soaks the sponge. Apply using repeated up and down, back and forth strokes for at least 30 seconds.



Dry

Leave the area to air **dry completely** before applying sterile drapes. Do not blot or wipe away. Discard the applicator after a single use.

Important safety point

Do not drape or use ignition source until the solution has completely dried.



1.5ml applicator



NHS list price per applicator:
£0.78 (ex. VAT and delivery)

NHS Supply chain
order code: MRB561

Coverage area:
10cm x 13cm

For procedures such as:

- Blood culture collection
- Peripheral canulation
- AV fistula and 'portacath' access
- Simple biopsies
- Venesection

ChloroPrep®
Patient Preoperative Skin Preparation
2% Chlorhexidine Gluconate (CHG) w/v
& 70% Isopropyl Alcohol (IPA) v/v

For customer services and all other enquiries:

Please telephone: 0800 043 7546
Email: enquiries@chloroprep.co.uk
or visit: www.chloroprep.co.uk



Prescribing Information ChloraPrep® (PL31760/0002) & ChloraPrep with Tint (PL31760-0001) 2% chlorhexidine gluconate w/v / 70% isopropyl alcohol v/v cutaneous solution. **Indication:** Disinfection of skin prior to invasive medical procedures. **Dosage & administration:** ChloraPrep – 0.67ml, 1.5ml, 3ml, 10.5ml, 26ml ; ChloraPrep with Tint – 3ml, 10.5ml, 26ml. Volume dependent on invasive procedure being undertaken. Applicator squeezed to break ampoule and release antiseptic solution onto sponge. Solution applied by gently pressing sponge against skin and moving back and forth for 30 seconds. The area covered should be allowed to air dry. **Side effects, precautions & contra-indications:** Very rarely allergic or skin reactions reported with chlorhexidine, isopropyl alcohol and Sunset Yellow. Contra-indicated for patients with known hypersensitivity to these constituents. For external use only on intact skin. Avoid contact with eyes, mucous membranes, middle ear and neural tissue. Should not be used in children under 2 months of age. Solution is flammable. Do not use with ignition sources until dry, do not allow to pool, and remove soaked materials before use. Over-vigorous use on fragile or sensitive skin or repeated use may lead to local skin reactions. At the first sign of local skin reaction, application should be stopped. **Per applicator costs (ex VAT)** ChloraPrep: 0.67ml (SEPP) - 30p; 1.5ml (FREPP) - 55p; 1.5ml – 78p; 3ml – 85p; 10.5ml - £2.92; 26ml - £6.50 ChloraPrep with Tint: 3ml – 89p; 10.5ml £3.07; 26ml - £6.83. **Legal category:** GSL. **Marketing Authorisation Holder:** CareFusion UK 244 Ltd, 43 London Road, Reigate, Surrey RH2 9PW, UK. **Date of preparation:** Oct 2010 CHLIH017