



We report a case where 2% chlorhexidine (CHX) gluconate was mistaken for an anesthetic solution and infiltrated into the buccal vestibule during routine root canal treatment” Hiremath et al (2016).

Abstract:

We report a case where 2% chlorhexidine (CHX) gluconate was mistaken for an anesthetic solution and infiltrated into the buccal vestibule during routine root canal treatment. Accidentally, 2% CHX gluconate solution was injected in the right upper buccal vestibule (16) of a 23-year-old male during routine root canal treatment. The patient experienced pain and a burning sensation over the injected area shortly after injection.

ReTweet if useful... Case study describes how 2% chlorhexidine was mistaken for local anaesthetic <http://ctt.ec/o7Fxb+> @ivteam #ivteam

Click To Tweet

Swelling with mild extraoral redness over the right cheek area was observed clinically. The patient was immediately administered dexamethasone intramuscularly, and was prescribed antibiotics, analgesics, and antihistamines. The patient complained of a loss of sensation over the right cheek by the 15(th) day. The swelling reduced gradually over a period of 15 days. Reversal of sensation was attained after 35 days.



Case study describes how 2% chlorhexidine was mistaken for local anaesthetic | 2

Full Text

Reference:

Hiremath, H., Agarwal, R.S., Patni, P. and Chauhan, S. (2016) Accidental injection of 2% chlorhexidine gluconate instead of an anesthetic agent: A case report. Journal of Conservative Dentistry. 19(1), p.106-8.

doi: 10.4103/0972-0707.173213.

Thank you to our partners for supporting IVTEAM

