

A peripherally inserted central catheter (PICC) was placed through the left basilic vein due to difficult intravenous access” Talari et al (2016).

Extract:

“A 31-year-old female patient with systolic heart failure (ejection fraction of 30%) and automatic implantable cardioverter-defibrillator (AICD) was admitted to the hospital with sepsis from *Staphylococcus hominis* and *Staphylococcus capitis* bacteraemia.

ReTweet if useful... Full text case study describe azygos vein PICC tip placement
[@ivteam #ivteam](http://ctt.ec/15130+)

Click To Tweet

A transesophageal echocardiogram showed AICD lead vegetation. Subsequently, the AICD was replaced. A peripherally inserted central catheter (PICC) was placed through the left basilic vein due to difficult intravenous access. Initially, fluoroscopy (figure 1) and chest X-ray (figure 2) confirmed appropriate position of the catheter tip in the superior vena cava (SVC)” Talari et al (2016).

[Full Text](#)

Reference:

Talari, G., Oyewole-Eletu, S., Talari, P. and Parasramka, S. (2016) Migration of peripherally inserted central catheter likely into the azygos vein: a conservative management. *BMJ Case Reports*. July 29th.

doi: 10.1136/bcr-2016-216657.

Thank you to our partners for supporting IVTEAM