

**This prompted insertion of a central venous catheter on the left side of the neck in the presence of normal vascular anatomy. A subsequent chest X-ray revealed an abnormal course of the catheter consistent with presence of persistent left superior vena cava”  
Sonawane et al (2017).**

Abstract:

Persistent left superior vena cava is a rare vascular anatomical variant. Although ultrasonography has facilitated the process of central venous catheterization, it cannot be used to locate the tip of a catheter. Postprocedure chest X-ray is desirable for locating the course and tip of the catheter. In our case, initial ultrasonography demonstrated the absence of the right internal jugular vein. This prompted insertion of a central venous catheter on the left side of the neck in the presence of normal vascular anatomy. A subsequent chest X-ray revealed an abnormal course of the catheter consistent with presence of persistent left superior vena cava.

ReTweet if useful... Persistent left superior vena cava and absent of right internal jugular vein [@ivteam #ivteam](https://ctt.ec/ag0ea+)

Click To Tweet

Reference:

Sonawane, N.B., Shah, F.R. and Gawankar, P.J. (2017) Accidental Cannulation of Persistent Left Superior Vena Cava in a Case of Absent Right Internal Jugular Vein: Role of Ultrasonography and X-Ray. A & A Case Reports. March 26th. .

doi: 10.1213/XAA.0000000000000505.

**Thank you to our partners for supporting IVTEAM**