

A peripherally inserted central catheter (PICC) is placed in the upper extremity with its distal end terminating in the superior vena cava. A midline catheter (MC) is a peripheral intravenous (IV) line measuring 3-8 inches and is inserted with its tip positioned just below the axilla” DiNucci et al (2016).

Abstract:

A peripherally inserted central catheter (PICC) is placed in the upper extremity with its distal end terminating in the superior vena cava. A midline catheter (MC) is a peripheral intravenous (IV) line measuring 3-8 inches and is inserted with its tip positioned just below the axilla. Both are utilized when long-term IV access is required or if venous access is limited. Our objective is to compare the utilization and safety of PICC lines versus midline catheters.

ReTweet if useful... Safety of peripherally inserted central catheters versus midline catheters
<http://ctt.ec/25nX6+> @ivteam #ivteam

Click To Tweet

Reference:

DiNucci, S., Messer, G., Xu, T., Shutt, K.A., Ferrelli, J., Andrzejewski, C. and Yassin, M.H. (2016) Comparison of Utilization and Safety of Peripherally Inserted Central Catheters Versus Midline Catheters in a Large Academic Medical Center. 44(6), Supplement, p.S71-S72.

DOI: <http://dx.doi.org/10.1016/j.ajic.2016.04.073>

Thank you to our partners for supporting IVTEAM