



We conducted this prospective study to evaluate the incidence of, major risk factors for, and causative pathogens of catheter-related bloodstream infection (CRBSI) related to central venous catheters (CVCs) in patients undergoing gastrointestinal surgery” Wu et al (2017).

Abstract:

We conducted this prospective study to evaluate the incidence of, major risk factors for, and causative pathogens of catheter-related bloodstream infection (CRBSI) related to central venous catheters (CVCs) in patients undergoing gastrointestinal surgery. The overall CRBSI rate was 8.0% (38/477), equaling 5.6 CRBSIs per 1,000 catheter days. CVC duration, emergency surgery, and male sex were risk factors for CRBSI. The most common organisms were coagulase-negative staphylococci, followed Enterobacteriaceae and Candida spp.

ReTweet if useful... What are the CLABSI risk factors after gastrointestinal surgery?
[@ivteam #ivteam](https://ctt.ec/036zg+)

Click To Tweet

Reference:

Wu, S., Ren, S., Zhao, H., Jin, H., Xv, L., Qian, S. and Wang, S. (2017) Risk factors for central venous catheter-related bloodstream infections after gastrointestinal surgery. American



What are the CLABSI risk factors after gastrointestinal surgery? | 2

Journal of Infection Control. February 20th. .

doi: 10.1016/j.ajic.2017.01.007.

Thank you to our partners for supporting IVTEAM

